



CO-PRODUCT BREEDER R1600 MINERAL GRA Medicated

A vitamin, mineral supplement for cattle (mature reproducing beef cows in dry lots); for improved feed efficiency.

ACTIVE DRUG INGREDIENT

Monensin Sodium 1600 g/ton

Guaranteed Analysis

Calcium (Ca)	(Min)	19.6 %	Copper (Cu)	(Min)	3,026 ppm
Calcium (Ca)	(Max)	23.5 %	Selenium (Se)	(Min)	25.0 ppm
Phosphorus (P)	(Min)	2.0 %	Zinc (Zn)	(Min)	4,029 ppm
Salt (NaCl)	(Min)	12.2 %	Vitamin A	(Min)	303,000 IU/lb
Salt (NaCl)	(Max)	14.6 %	Vitamin D-3	(Min)	30,300 IU/lb
Magnesium (Mg)	(Min)	2.1 %	Vitamin E	(Min)	308 IU/lb
Potassium (K)	(Min)	1.1 %	Thiamin	(Min)	814 mg/lb

List of Ingredients

Calcium Carbonate, Distillers Dried Grains/Solubles, Salt, Dicalcium Phosphate, Magnesium Oxide, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Potassium Chloride, Sodium Selenite, Cane Molasses, Mineral Oil, Ethylenediamine Dihydroiodide, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ethoxyquin Preservative, Sodium Sesquicarbonate, Yeast Culture, Thiamine Mononitrate, Natural and Artificial Flavor, Choline Chloride, Niacin, Calcium Pantothenate, Riboflavin, Vitamin B-12 Supplement, Biotin, Menadione Dimethylpyrimidinol Bisulfite (Source of Vitamin K Activity), Thiamine, Pyridoxine Hydrochloride, Folic Acid, Artificial Color.

Feeding Directions

Mix 3 oz. CO-PRODUCT BREEDER R1600 MINERAL GRA with 13 oz. of grain and hand feed at the rate of 1 lb. per head, per day to provide 150 mg of Monensin. During the first five days of feeding, cattle should receive no more than 100 mg per day contained in not less than 1 pound of feed.

Caution

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle and goats only. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. Due to the copper level, do not feed to sheep. Follow label directions. The feeding of supplemental selenium should not exceed 0.3 ppm of total diet. The addition to feed of higher levels of this premix containing selenium is not permitted.

WARNING: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

Manufactured by:



6010 Stockyards Expressway, St. Joseph, MO 64504
(800) 821-3070

Net Weight 50 lbs. (22.7 kg)

31 10/28/2016 111034W

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed. Safety Data Sheet available upon request.