



CATTLE DEVELOPER RU1200 GRAN MEDICATED

A vitamin and mineral supplement for beef cattle fed in confinement for slaughter for improved feed efficiency

ACTIVE DRUG INGREDIENT

Monensin (as monensin sodium)..... 1200 g/ton

*** Contains not more than 8.5 % protein from non-protein nitrogen.**

Guaranteed Analysis

Crude Protein*	(Min)	10.4 %	Potassium (K)	(Min)	2.5 %
Crude Fat	(Min)	1.6 %	Copper (Cu)	(Min)	2,003 ppm
Crude Fiber	(Max)	0.7 %	Selenium (Se)	(Min)	20.0 ppm
Calcium (Ca)	(Min)	22.8 %	Vitamin A	(Min)	302,600 IU/lb
Calcium (Ca)	(Max)	27.3 %	Vitamin D-3	(Min)	30,700 IU/lb
Phosphorus (P)	(Min)	3.0 %	Vitamin E	(Min)	303 IU/lb
Salt (NaCl)	(Min)	5.3 %	Thiamin	(Min)	167 mg/lb
Salt (NaCl)	(Max)	6.3 %			

List of Ingredients

Calcium Carbonate, Dicalcium Phosphate, Distillers Dried Grains/Solubles, Salt, Potassium Chloride, Urea, Magnesium Oxide, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Ethylenediamine Dihydroiodide, Mineral Oil, Cobalt Sulfate, Calcium Iodate, Dried Aspergillus oryzae Fermentation Product, Sodium Selenite, Cane Molasses, Soybean Oil, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ethoxyquin Preservative, Yeast Culture, Natural and Artificial Flavor, Thiamine Mononitrate, Artificial Color, Choline Chloride, Niacin, Calcium Pantothenate, Riboflavin, Vitamin B-12 Supplement, Biotin, Menadione Dimethylpyrimidinol Bisulfite (Source of Vitamin K Activity), Thiamine, Pyridoxine Hydrochloride, Folic Acid.

Contains AMAFERM®, a registered trademark of BioZyme® Inc

Feeding Directions

Feed 1/4-1/3 lb. per head, per day mixed in a grain mix or TMR to provide 150 to 200 mg of Monensin daily. Do not free-choice.

Caution

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle and goats only. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. Due to the copper level, do not feed to sheep. Follow label directions. The feeding of supplemental selenium should not exceed 0.3 ppm of total diet. The addition to feed of higher levels of this premix containing selenium is not permitted. WARNING: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

Manufactured by:



6010 Stockyards Expressway, St. Joseph, MO 64504
(800) 821-3070

Net Weight 50 lbs. (22.7 kg)

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed. Safety Data Sheet available upon request.

31 10/18/2016 111100