



SUP'R PIG 15-25 DEN35/AUREO400 MP Medicated

CAUTION: Beginning January 1, 2017, Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For control of swine dysentery associated with Brachyspira (formerly Serpulina or Treponema) hyodysenteriae susceptible to Tiamulin. Treatment of bacterial enteritis caused by Escherichia coli and Salmonella choleraesuis and bacterial pneumonia caused by Pasteurella multocida susceptible to Chlortetracycline.

ACTIVE DRUG INGREDIENTS

Tiamulin hydrogen fumarate	35 g/ton
Chlortetracycline	400 g/ton

Guaranteed Analysis

Crude Protein	(Min)	20.7 %	Calcium (Ca)	(Max)	1.2 %
Lysine	(Min)	1.37 %	Phosphorus (P)	(Min)	0.76 %
Crude Fat	(Min)	6.0 %	Selenium (Se)	(Min)	0.3 ppm
Crude Fiber	(Max)	2.3 %	Zinc (Zn)	(Min)	2,667 ppm
Calcium (Ca)	(Min)	0.7 %	Vitamin A	(Min)	6,900 IU/lb

List of Ingredients

Ground Yellow Corn, Soybean Meal, Dried Whey, Oat Groats, Fish Meal, Choice White Grease, Lactose, Dicalcium Phosphate, Calcium Carbonate, Zinc Oxide, Amino Resin Calcium Sulfate, Artificial Flavor, L-Lysine, Sodium Selenite, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Choline Chloride, Niacin, Calcium Pantothenate, Riboflavin, Vitamin B-12 Supplement, Biotin, Menadione Dimethylpyrimidinol Bisulfite (Source of Vitamin K Activity), Thiamine, Pyridoxine Hydrochloride, Folic Acid, Ethoxyquin Preservative, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Ethylenediamine Dihydroiodide, Mineral Oil, DL-Methionine, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide.

Feeding Directions

SUP'R PIG 15-25 DEN35G/AUREO400G MP Medicated is designed to be fed continuously as the sole ration for 14 days to pigs from 15-25 lbs. bodyweight.

Caution

CAUTION: Do not feed to animals other than swine. Withdraw 2 days before slaughter. SPECIAL CONSIDERATION: Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g., Monensin, Lasalocid, Narasin, Salinomycin, and Semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use. The effects of Tiamulin on swine reproductive performance, pregnancy, and lactation have not been determined.

LIMITATIONS: Feed approximately 400 g/ton of Chlortetracycline, varying with bodyweight and feed consumption to provide 10 mg/lb. of bodyweight per day. Feed continuously as sole ration for 14 days. Use as only source of chlortetracycline. Use as only source of tiamulin. The feeding of supplemental selenium should not exceed 0.3 ppm of total diet. The addition to feed of higher levels of this premix containing selenium is not permitted.

Manufactured by:



6010 Stockyards Expressway, St. Joseph, MO 64504
(800) 821-3070

Net Weight 50 lbs. (22.7 kg)

31 12/08/2016 129715A

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed. Safety Data Sheet available upon request.