

TOP PIG 16% T40 MEAL

Medicated

Sole ration for swine from 120 lb. to market weight for control of swine dysentery caused by Brachyspira hyodysenteriae; and for control of porcine proliferative enteropathies (ileitis) associated with Lawsonia intracellularis.

CAUTION: Beginning January 1, 2017, Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

ACTIVE DRUG INGREDIENT

ACTIVE BROOTHOREBIET					
	Tylosin			40 g/ton	
Guaranteed Analysis					
Crude Protein	(Min)	16.1 %	Salt (NaCl)	(Min)	0.10 %
Lysine	(Min)	1.02 %	Salt (NaCl)	(Max)	0.60 %
Crude Fat	(Min)	5.4 %	Selenium (Se)	(Min)	0.3 ppm
Crude Fiber	(Max)	2.7 %	Zinc (Zn)	(Min)	174 ppm
Calcium (Ca)	(Min)	0.5 %	Vitamin A	(Min)	4,700 IU/lb
Calcium (Ca)	(Max)	1.0 %	Vitamin D-3	(Min)	600 IU/lb
Phosphorus (P)	(Min)	0.60 %	Vitamin E	(Min)	26 IU/lb
List of Ingredients					

Ground Yellow Corn, Soybean Meal, Oat Groats, Choice White Grease, Dicalcium Phosphate, Dried Whey, Calcium Carbonate, Animal Fat with BHA preservative, Lecithin, Casein, Fish Meal, L-Lysine, Lactose, Monocalcium Phosphate, Salt, Sodium Selenite, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Ethylenediamine Dihydroiodide, Mineral Oil, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Choline Chloride, Niacin, Calcium Pantothenate, Riboflavin, Vitamin B-12 Supplement, Biotin, Menadione Dimethylpyrimidinol Bisulfite (Source of Vitamin K Activity), Thiamine, Pyridoxine Hydrochloride, Folic Acid, Ethoxyquin Preservative, Artificial Flavor, DL-Methionine, Zinc Oxide.

Feeding Directions

Feed continuously as the sole ration to growing swine. Provide clean, fresh water at all times.

Caution

NOTICE: Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of $2\%\,$

LIMITATIONS: Include 100 g/ton Tylosin for at least three weeks, followed by 40 g/ton until market weight.

CAUTION: The feeding of supplemental selenium should not exceed 0.3 ppm of total diet. The addition to feed of higher levels of this premix containing selenium is not permitted.

Manufactured by:





6010 Stockyards Expressway, St. Joseph, MO 64504 (800) 821-3070

Net Weight 50 lbs. (22.7 kg)

31 07/13/2016 13432X

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed. Safety Data Sheet available upon request.