



FDLT RECEIVER 5 OXY800 PEL MEDICATED

A protein, vitamin, and mineral supplement for weaning and receiving beef cattle for the prevention and treatment of the early stages of shipping fever complex. **CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

ACTIVE DRUG INGREDIENT

Oxytetracycline 800 g/ton

Guaranteed Analysis

Crude Protein	(Min)	15.0 %	Salt (NaCl)	(Max)	0.9 %
Crude Fat	(Min)	4.4 %	Potassium (K)	(Min)	1.1 %
Crude Fiber	(Max)	17.4 %	Selenium (Se)	(Min)	1.3 ppm
Calcium (Ca)	(Min)	1.5 %	Vitamin A	(Min)	33,900 IU/lb
Calcium (Ca)	(Max)	2.0 %	Vitamin D-3	(Min)	3,400 IU/lb
Phosphorus (P)	(Min)	0.60 %	Vitamin E	(Min)	88 IU/lb
Salt (NaCl)	(Min)	0.4 %	Niacin	(Min)	502 mg/lb

List of Ingredients

Wheat Middlings, Soybean Hulls, Ground Yellow Corn, Linseed Meal, Distillers Dried Grains/Solubles, Calcium Carbonate, Cane Molasses, Yeast Culture, Soybean Meal, Choice White Grease, Lignin Sulfonate, Pork Meat & Bone Meal, Salt, Potassium Chloride, Vitamin E Supplement, Niacin, Zinc Proteinate, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Dried Aspergillus oryzae Fermentation Product, Manganese Proteinate, Sodium Selenite, Copper Proteinate, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide, Vitamin A Supplement, Vitamin D-3 Supplement, Choline Chloride, Calcium Pantothenate, Riboflavin, Vitamin B-12 Supplement, Biotin, Menadione Dimethylpyrimidinol Bisulfite (Source of Vitamin K Activity), Thiamine, Pyridoxine Hydrochloride, Folic Acid, Ethoxyquin Preservative, Thiamine Mononitrate, Cobalt Carbonate.

Contains Amaferm®, a registered trademark of BioZyme® Inc.

Feeding Directions

Feed FDLT RECEIVER 5 OXY800 PEL MEDICATED at the rate of 5 lbs. per head, per day to newly weaned or received beef calves 3-5 days preceding shipment and 3-5 days following arrival in feedlots. FDLT RECEIVER 5 OXY800 PEL MEDICATED provides 2 grams of oxytetracycline daily.

Caution

SPECIAL CONSIDERATIONS: For use in dry animal feed only. Not for use in liquid feed supplements. The feeding of supplemental selenium should not exceed 0.3 ppm of total diet. The addition to feed of higher levels of this premix containing selenium is not permitted.

Manufactured by:



6010 Stockyards Expressway, St. Joseph, MO 64504
(800) 821-3070

Net Weight 50 lbs. (22.7 kg)

31 08/01/2018 143600

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed. Safety Data Sheet available upon request.