

# FDLT RECEIVER 2 MED PEL Medicated

A vitamin and mineral supplement for beef cattle fed in confinement for slaughter for the prevention and treatment of the early stages of shipping fever complex. CAUTION: Federal law restricts medicated feed containing this veterinary feed

directive (VFD) drug to use by or on the order of a licensed veterinarian.

### ACTIVE DRUG INGREDIENT

Oxytetracycline ..... 1 g/lb

Guaranteed Analysis					
Crude Protein	(Min)	24.1 %	Potassium (K)	(Min)	2.0 %
Crude Fat	(Min)	4.1 %	Selenium (Se)	(Min)	3.0 ppm
Crude Fiber	(Max)	8.3 %	Vitamin A	(Min)	75,800 IU/lb
Calcium (Ca)	(Min)	2.0 %	Vitamin D-3	(Min)	7,600 IU/lb
Calcium (Ca)	(Max)	2.5 %	Vitamin E	(Min)	200 IU/lb
Phosphorus (P)	(Min)	0.83 %	Niacin	(Min)	1,250 mg/lb
Salt (NaCl)	(Min)	1.0 %	Thiamine	(Min)	45 mg/lb
Salt (NaCl)	(Max)	1.5 %			
List of Ingradiante					

#### **List of Ingredients**

Distillers Dried Grains/Solubles, Linseed Meal, Wheat Middlings, Soybean Meal, Yeast Culture, Cane Molasses, Calcium Carbonate, Pork Meat & Bone Meal, Potassium Chloride, Soybean Hulls, Salt, Vitamin E Supplement, Niacin, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Dicalcium Phosphate, Dried Aspergillus oryzae Fermentation Product, Sodium Selenite, Zinc Proteinate, Vitamin A Supplement, Vitamin D-3 Supplement, Manganese Proteinate, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide, Choline Chloride, Calcium Pantothenate, Riboflavin, Vitamin B-12 Supplement, Biotin, Menadione Dimethylpyrimidinol Bisulfite (Source of Vitamin K Activity), Thiamine, Pyridoxine Hydrochloride, Folic Acid, Ethoxyquin Preservative, Copper Proteinate, Thiamine Mononitrate.

Contains AMAFERM®, a registered trademark of BioZyme® Inc.

#### **Feeding Directions**

Feed FDLT RECEIVER 2 MED PEL Medicated at the rate of 2 lbs. per head, per day to newly weaned or received beef calves to provide 2 g of Oxytetracycline daily.

Caution

SPECIAL CONSIDERATIONS: For use in dry animal feed only. Not for use in liquid feed supplements. The feeding of supplemental selenium should not exceed 0.3 ppm of total diet. The addition to feed of higher levels of this premix containing selenium is not permitted.

## Manufactured by:



MEDICATED

6010 Stockyards Expressway, St. Joseph, MO 64504 (800) 821-3070 Net Weight 50 lbs. (22.7 kg) 31 03/09/2018 145600

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed. Safety Data Sheet available upon request.