



FDLT NATURAL RU400 PEL Medicated

A vitamin and mineral supplement for beef cattle fed in confinement for slaughter for improved feed efficiency

ACTIVE DRUG INGREDIENT

Monensin (as monensin sodium)..... 400.0 g/ton

Guaranteed Analysis

Crude Protein	(Min)	32.0 %	Salt (NaCl)	(Max)	2.7 %
Crude Fat	(Min)	3.2 %	Potassium (K)	(Min)	1.5 %
Crude Fiber	(Max)	4.6 %	Selenium (Se)	(Min)	2.2 ppm
Calcium (Ca)	(Min)	6.6 %	Vitamin A	(Min)	41,200 IU/lb
Calcium (Ca)	(Max)	8.0 %	Vitamin D-3	(Min)	4,100 IU/lb
Phosphorus (P)	(Min)	0.86 %	Vitamin E	(Min)	42 IU/lb
Salt (NaCl)	(Min)	2.2 %	Thiamin	(Min)	105 mg/lb

List of Ingredients

Soybean Meal, Distillers Dried Grains/Solubles, Calcium Carbonate, Pork Meat & Bone Meal, Wheat Middlings, Cane Molasses, Salt, Linseed Meal, Limestone ground, Lignin Sulfonate, Dried Aspergillus oryzae Fermentation Product, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Dehydrated Alfalfa Meal, Potassium Chloride, Sodium Selenite, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ethoxyquin Preservative, Mineral Oil, Thiamine Mononitrate.

Contains AMAFERM®, a registered trademark of BioZyme® Inc.

Feeding Directions

Feed 1 pound of FDLT NATURAL RU400 PEL Medicated per head per day to feedlot cattle to provide 200 mg of Monensin daily.

Caution

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle and goats only. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. The feeding of supplemental selenium should not exceed 0.3 ppm of total diet. The addition to feed of higher levels of this premix containing selenium is not permitted.

WARNING: A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal.

Manufactured by:



MEDICATED

6010 Stockyards Expressway, St. Joseph, MO 64504
(800) 821-3070

Net Weight 50 lbs. (22.7 kg)

31 03/16/2017 145934E

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed. Safety Data Sheet available upon request.