



FDLT 40/14 BT500 M.4 PEL Medicated

A protein, vitamin and mineral supplement for increased rate of weight gain, improved feed efficiency and suppression of estrus (heat) in heifers fed in confinement for slaughter.

ACTIVE DRUG INGREDIENTS

Melengestrol Acetate	0.4 mg/lb
Lasalocid Sodium	500 g/ton

*** Contains not more than 14.0 % protein from non-protein nitrogen.**

Guaranteed Analysis

Crude Protein*	(Min)	40.1 %	Salt (NaCl)	(Max)	3.1 %
Crude Fat	(Min)	2.3 %	Potassium (K)	(Min)	1.1 %
Crude Fiber	(Max)	4.0 %	Selenium (Se)	(Min)	3.2 ppm
Calcium (Ca)	(Min)	10.4 %	Vitamin A	(Min)	41,200 IU/lb
Calcium (Ca)	(Max)	12.4 %	Vitamin D-3	(Min)	4,100 IU/lb
Phosphorus (P)	(Min)	0.70 %	Vitamin E	(Min)	42 IU/lb
Salt (NaCl)	(Min)	2.6 %			

List of Ingredients

Soybean Meal, Calcium Carbonate, Linseed Meal, Distillers Dried Grains/Solubles, Pork Meat & Bone Meal, Urea, Cane Molasses, Salt, Limestone ground, Magnesium Oxide, Dehydrated Alfalfa Meal, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Sodium Selenite, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement.

Feeding Directions

Feed FDLT 40/14 BT500 M.4 PEL Medicated continuously at the rate of 1 pound per head per day to heifers being fed in confinement for slaughter to provide 250 mg of Lasalocid and .4 mg Melengestrol Acetate daily.

Consult your PEET'S Representative for ration balancing services and medicated options available.

NOTICE: Expiration date of 90 days after manufacture.

Caution

Not effective in steers and spayed heifers. The safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to supplements containing lasalocid as ingestion may be fatal. Mix thoroughly with grain and(or) roughage prior to feeding. Feeding undiluted, mixing errors and inadequate mixing of the total ration may result in excess lasalocid concentration which could be fatal to cattle. A withdrawal period has not been established for lasalocid in pre-ruminating cattle. Do not use in calves to be processed for veal. The feeding of supplemental selenium should not exceed 0.3 ppm of total diet. The addition to feed of higher levels of this premix containing selenium is not permitted.

Manufactured by:



MEDICATED

6010 Stockyards Expressway, St. Joseph, MO 64504
(800) 821-3070

Net Weight 50 lbs. (22.7 kg)

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed. Safety Data Sheet available upon request.

31 07/11/2017 146627E