



# FDLT 40/14 RU400 PEL Medicated

A protein, vitamin and mineral supplement for beef cattle fed in confinement for slaughter for improved feed efficiency

### ACTIVE DRUG INGREDIENT

Monensin (as monensin sodium)..... 400.0 g/ton

**\* Contains not more than 13.9 % protein from non-protein nitrogen.**

### Guaranteed Analysis

Crude Protein*	(Min)	40.1 %	Salt (NaCl)	(Max)	3.1 %
Crude Fat	(Min)	2.5 %	Potassium (K)	(Min)	1.1 %
Crude Fiber	(Max)	4.1 %	Selenium (Se)	(Min)	3.2 ppm
Calcium (Ca)	(Min)	10.3 %	Vitamin A	(Min)	41,200 IU/lb
Calcium (Ca)	(Max)	12.4 %	Vitamin D-3	(Min)	4,100 IU/lb
Phosphorus (P)	(Min)	0.71 %	Vitamin E	(Min)	42 IU/lb
Salt (NaCl)	(Min)	2.6 %			

### List of Ingredients

Soybean Meal, Calcium Carbonate, Distillers Dried Grains/Solubles, Linseed Meal, Pork Meat & Bone Meal, Urea, Cane Molasses, Salt, Limestone ground, Dehydrated Alfalfa Meal, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Sodium Selenite, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement.

### Feeding Directions

Feed 1 pound of FDLT 40/14 RU400 PEL Medicated per head per day to feedlot cattle to provide 200 mg of Monensin daily.

### Caution

**CAUTION:** Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle and goats only. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. The feeding of supplemental selenium should not exceed 0.3 ppm of total diet. The addition to feed of higher levels of this premix containing selenium is not permitted.

**WARNING:** A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal.

**LIMITATIONS:** Feed continuously in complete feed at a rate of 50 to 480 milligrams of monensin per head per day. No additional improvement in feed efficiency has been shown from feeding monensin at levels greater than 30 grams per ton.

Manufactured by:



**MEDICATED**

6010 Stockyards Expressway, St. Joseph, MO 64504

(800) 821-3070

**Net Weight 50 lbs. (22.7 kg)**

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed. Safety Data Sheet available upon request.

31 07/21/2017 146634E