



# FEEDLOT PERFORMER RU1200 PEL

## Medicated

A vitamin and mineral supplement for beef cattle fed in confinement for slaughter: For improved feed efficiency.

### ACTIVE DRUG INGREDIENT

Monensin Sodium ..... 1200 g/ton

### Guaranteed Analysis

|                |       |        |               |       |               |
|----------------|-------|--------|---------------|-------|---------------|
| Crude Protein  | (Min) | 10.0 % | Salt (NaCl)   | (Max) | 5.1 %         |
| Crude Fat      | (Min) | 2.9 %  | Potassium (K) | (Min) | 2.6 %         |
| Crude Fiber    | (Max) | 8.4 %  | Selenium (Se) | (Min) | 6.6 ppm       |
| Calcium (Ca)   | (Min) | 10.5 % | Zinc (Zn)     | (Min) | 1,030 ppm     |
| Calcium (Ca)   | (Max) | 12.5 % | Vitamin A     | (Min) | 160,800 IU/lb |
| Phosphorus (P) | (Min) | 2.00 % | Vitamin D-3   | (Min) | 16,000 IU/lb  |
| Salt (NaCl)    | (Min) | 4.1 %  | Vitamin E     | (Min) | 164 IU/lb     |

### List of Ingredients

Wheat Middlings, Calcium Carbonate, Distillers Dried Grains/Solubles, Dicalcium Phosphate, Soybean Hulls, Salt, Potassium Chloride, Cane Molasses, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Choice White Grease, Sodium Selenite, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ethoxyquin Preservative, Mineral Oil, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide.

### Feeding Directions

Feed FEEDLOT PERFORMER RU1200 PEL Medicated continuously to cattle being fed in confinement for slaughter at the rate of 0.33 to 0.50 lb. per head, per day, to provide 200 to 300 mg of Monensin.

### Caution

**CAUTION:** Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle and goats only. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. The feeding of supplemental selenium should not exceed 0.3 ppm of total diet. The addition to feed of higher levels of this premix containing selenium is not permitted.

**WARNING:** A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal.

Manufactured by:



6010 Stockyards Expressway, St. Joseph, MO 64504  
(800) 821-3070

**Net Weight 50 lbs. (22.7 kg)**

**31 03/20/2017 149334H**

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed. Safety Data Sheet available upon request.