



FEEDLOT CP BALANCER R1000 GRAN Medicated

A vitamin and mineral supplement for beef cattle fed in confinement for slaughter for improved feed efficiency

ACTIVE DRUG INGREDIENT

Monensin (as monensin sodium)..... 1000 g/ton

Guaranteed Analysis

Calcium (Ca)	(Min)	28.5 %	Selenium (Se)	(Min)	8.8 ppm
Calcium (Ca)	(Max)	33.5 %	Zinc (Zn)	(Min)	2,009 ppm
Salt (NaCl)	(Min)	12.4 %	Vitamin A	(Min)	103,100 IU/lb
Salt (NaCl)	(Max)	14.9 %	Vitamin D-3	(Min)	10,300 IU/lb
Potassium (K)	(Min)	2.0 %	Vitamin E	(Min)	103 IU/lb
Copper (Cu)	(Min)	502 ppm	Thiamine	(Min)	104 mg/lb

List of Ingredients

Calcium Carbonate, Salt, Potassium Chloride, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Wheat Middlings, Mineral Oil, Sodium Selenite, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Thiamine Mononitrate.

Feeding Directions

Mix and feed 1/2 pound of FEEDLOT CP BALANCER R1000 GRAN Medicated daily to beef cattle being fed in confinement for slaughter to provide 250 mg of Monensin.

Caution

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle and goats only. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. Due to the copper level, do not feed to sheep. Follow label directions. The feeding of supplemental selenium should not exceed 0.3 ppm of total diet. The addition to feed of higher levels of this premix containing selenium is not permitted.

WARNING: A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal.

LIMITATIONS: Feed continuously in complete feed at a rate of 50 to 480 milligrams of monensin per head per day. No additional improvement in feed efficiency has been shown from feeding monensin at levels greater than 30 grams per ton.

Manufactured by:



6010 Stockyards Expressway, St. Joseph, MO 64504
(800) 821-3070

Net Weight 50 lbs. (22.7 kg)

31 05/03/2017 149634G

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed. Safety Data Sheet available upon request.