



# FDLT CP BALANCER RU1200 GRAN Medicated

A vitamin and mineral supplement for beef cattle fed in confinement for slaughter for improved feed efficiency

### ACTIVE DRUG INGREDIENT

Monensin (as monensin sodium)..... 1200.0 g/ton

### Guaranteed Analysis

Calcium (Ca)	(Min)	28.0 %	Selenium (Se)	(Min)	8.8 ppm
Calcium (Ca)	(Max)	33.0 %	Zinc (Zn)	(Min)	3,000 ppm
Salt (NaCl)	(Min)	12.2 %	Vitamin A	(Min)	123,700 IU/lb
Salt (NaCl)	(Max)	14.7 %	Vitamin D-3	(Min)	12,300 IU/lb
Potassium (K)	(Min)	3.0 %	Vitamin E	(Min)	124 IU/lb
Copper (Cu)	(Min)	750 ppm	Thiamine	(Min)	209 mg/lb

### List of Ingredients

Calcium Carbonate, Salt, Potassium Chloride, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Mineral Oil, Sodium Selenite, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Thiamine Mononitrate.

### Feeding Directions

Feed 1/4-1/3 lb. per head, per day mixed in a grain mix or TMR to provide 150 to 200 mg of Monensin daily. Do not free-choice.

### Caution

**CAUTION:** Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle and goats only. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. Due to the copper level, do not feed to sheep. Follow label directions. The feeding of supplemental selenium should not exceed 0.3 ppm of total diet. The addition to feed of higher levels of this premix containing selenium is not permitted.

**WARNING:** A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal.

**LIMITATIONS:** Feed continuously in complete feed at a rate of 50 to 480 milligrams of monensin per head per day. No additional improvement in feed efficiency has been shown from feeding monensin at levels greater than 30 grams per ton.

Manufactured by:



**MEDICATED**

6010 Stockyards Expressway, St. Joseph, MO 64504

(800) 821-3070

**Net Weight 50 lbs. (22.7 kg)**

**31 05/03/2017 149634H**

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed. Safety Data Sheet available upon request.