



TOP LAMB 32 D180 MP

Medicated

For the prevention of coccidiosis in young sheep caused by Eimeria ovinoidalis, E. crandallis, E. parva and E. bakuensis.

ACTIVE DRUG INGREDIENT

Decoquinatate 180 g/ton

*** Contains not more than 3.3 % protein from non-protein nitrogen.**

Guaranteed Analysis

Crude Protein*	(Min)	32.1 %	Salt (NaCl)	(Max)	4.4 %
Crude Fat	(Min)	3.4 %	Potassium (K)	(Min)	1.6 %
Crude Fiber	(Max)	7.4 %	Selenium (Se)	(Min)	1.2 ppm
Calcium (Ca)	(Min)	3.3 %	Zinc (Zn)	(Min)	1,122 ppm
Calcium (Ca)	(Max)	4.3 %	Vitamin A	(Min)	41,200 IU/lb
Phosphorus (P)	(Min)	0.89 %	Vitamin D-3	(Min)	4,100 IU/lb
Salt (NaCl)	(Min)	3.4 %	Vitamin E	(Min)	103 IU/lb

List of Ingredients

Soybean Meal, Distillers Dried Grains/Solubles, Wheat Middlings, Pork Meat & Bone Meal, Calcium Carbonate, Soybean Hulls, Linseed Meal, Cane Molasses, Salt, Ammonium Chloride, Potassium Sulfate, Magnesium Sulfate, Sodium Molybdenate, Lignin Sulfonate, Vitamin E Supplement, Zinc Sulfate, Zinc Oxide, Sodium Selenite, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide, Vitamin A Supplement, Vitamin D-3 Supplement, Ethylenediamine Dihydroiodide.

Feeding Directions

Mix Top Lamb 32 D180 with grain of choice as directed below and feed at a rate to provide 22.7 mg per 100 lb. of bodyweight of Decoquinatate daily for the prevention of coccidiosis in young sheep. Offer good quality hay and fresh clean water. Feed for at least 28 days during periods of exposure to coccidiosis or when experience indicates that it is likely to be a hazard.

	14% CP	13% CP
	Growing	Finishing
	50-100 lb.	100 lb.-MKT
Top Lamb 32	500	400
Grain (whole)	1,500	1,600
total	2,000	2,000
Decoquinatate	45 g/t	36 g/t

Caution

WARNING: Do not feed to sheep producing milk for food.

CAUTION: Do not use in feeds containing bentonite. The feeding of supplemental selenium should not exceed 0.3 ppm of total diet. The addition to feed of higher levels of this premix containing selenium is not permitted. Use only as directed. For ruminants (cattle, sheep and goats) only.

LIMITATIONS: Feed at least 28 days during periods of exposure to coccidiosis or when experience indicates that it is likely to be a hazard. Feed Type C medicated feed supplements as a top dress or mix into the daily ration to provide 22.7 mg decoquinatate per 100 lb (0.5 mg/kg) bodyweight. Type C feed may be prepared from dry or liquid type B feed.

Manufactured by:



MEDICATED

6010 Stockyards Expressway, St. Joseph, MO 64504
(800) 821-3070

Net Weight 50 lbs. (22.7 kg)

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed. Safety Data Sheet available upon request.

31 08/31/2017 157400