



MGA PAC 1 PEL Medicated

A vitamin, mineral supplement for increased rate of weight gain, improved feed efficiency and suppression of estrus (heat) in heifers fed for slaughter.

ACTIVE DRUG INGREDIENT

Melengestrol Acetate 1 mg/lb

Guaranteed Analysis

Crude Protein	(Min)	7.9 %	Phosphorus (P)	(Min)	0.39 %
Crude Fat	(Min)	2.8 %	Salt (NaCl)	(Min)	4.2 %
Crude Fiber	(Max)	12.5 %	Salt (NaCl)	(Max)	5.2 %
Calcium (Ca)	(Min)	14.3 %	Potassium (K)	(Min)	0.6 %
Calcium (Ca)	(Max)	17.1 %			

List of Ingredients

Calcium Carbonate, Wheat Middlings, Soybean Hulls, Salt, Cane Molasses, Choice White Grease.

Feeding Directions

Feed MGA PAC 1 PEL to heifers at the rate of 1/2 lb. per head, per day to provide 0.5 mg Melengestrol Acetate daily. Consult your PEET'S Representative for ration balancing services to provide the optimum usage of grain and roughage.

Caution

Feed MGA PAC 1 PEL to heifers at the rate of 1/2 lb. per head, per day to provide 0.5 mg Melengestrol Acetate daily. Consult your PEET'S Representative for ration balancing services to provide the optimum usage of grain and roughage.

Caution

CAUTION: Withdrawal periods of 3 to 5 days should be avoided to prevent the possibility that heifers may come into estrus (heat) at loading time. Not effective in steers and spayed heifers.

CAUTION: Heifers intended for reproduction: Do not exceed 24 days of feeding. A reduced conception rate can be expected if heifers are bred at estruses observed within 1 to 12 days after withdrawal of Melengestrol Acetate, whereas heifers bred at subsequent observed estrus are expected to have normal conception rates.

Manufactured by:

MEDICATED



6010 Stockyards Expressway, St. Joseph, MO 64504
(800) 821-3070

Net Weight 50 lbs. (22.7 kg)

31 05/05/2017 17202M

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed. Safety Data Sheet available upon request.