

TOP PIG 20% T40 MEAL

Medicated

Sole ration for swine from 120 lb. to market weight for control of swine dysentery caused by Brachyspira hyodysenteriae; and for control of porcine proliferative enteropathies (ileitis) associated with Lawsonia intracellularis.

CAUTION: Beginning January 1, 2017, Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

ACTIVE DRUG INGREDIENT

| | Tylosin | | | 40 g/ton | |
|---------------------|---------|--------|---------------|----------|-------------|
| Guaranteed Analysis | | | | | |
| Crude Protein | (Min) | 20.1 % | Salt (NaCl) | (Min) | 0.10 % |
| Lysine | (Min) | 1.20 % | Salt (NaCl) | (Max) | 0.60 % |
| Crude Fat | (Min) | 4.3 % | Selenium (Se) | (Min) | 0.3 ppm |
| Crude Fiber | (Max) | 2.9 % | Zinc (Zn) | (Min) | 138 ppm |
| Calcium (Ca) | (Min) | 0.6 % | Vitamin A | (Min) | 4,700 IU/lb |
| Calcium (Ca) | (Max) | 1.1 % | Vitamin D-3 | (Min) | 600 IU/lb |
| Phosphorus (P) | (Min) | 0.66 % | Vitamin E | (Min) | 25 IU/lb |
| List of Ingredients | | | | | |

Ground Yellow Corn, Soybean Meal, Dicalcium Phosphate, Choice White Grease, Calcium Carbonate, Dried Whey, Fish Meal, Animal Fat with BHA preservative, Lecithin, Casein, Monocalcium Phosphate, Salt, L-Lysine, Sodium Selenite, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide, Lactose, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Ethylenediamine Dihydroiodide, Mineral Oil, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Choline Chloride, Niacin, Calcium Pantothenate, Riboflavin, Vitamin B-12 Supplement,

Biotin, Menadione Dimethylpyrimidinol Bisulfite (Source of Vitamin K Activity), Thiamine, Pyridoxine Hydrochloride, Folic Acid, Ethoxyquin Preservative, Artificial Flavor, DL-Methionine.

Feeding Directions

Feed continuously as the sole ration to growing swine. Provide clean, fresh water at all times.

Caution

NOTICE: Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of $2\%\,$

LIMITATIONS: Include 100 g/ton Tylosin for at least three weeks, followed by 40 g/ton until market weight.

The feeding of supplemental selenium should not exceed 0.3 ppm of total diet. The addition to feed of higher levels of this premix containing selenium is not permitted.

Manufactured by:





6010 Stockyards Expressway, St. Joseph, MO 64504 (800) 821-3070

Net Weight 50 lbs. (22.7 kg)

31 07/13/2016 13432W

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed. Safety Data Sheet available upon request.