



FDLT NATURAL BT500 M.4 PEL

Medicated

For increased rate of weight gain, improved feed efficiency and suppression of estrus (heat) in heifers fed in confinement for slaughter.

ACTIVE DRUG INGREDIENTS

Melengestrol Acetate	0.4 mg/lb
Lasalocid Sodium	500 g/ton

Guaranteed Analysis

Crude Protein	(Min)	32.0 %	Salt (NaCl)	(Max)	3.2 %
Crude Fat	(Min)	3.2 %	Potassium (K)	(Min)	1.5 %
Crude Fiber	(Max)	3.9 %	Selenium (Se)	(Min)	2.2 ppm
Calcium (Ca)	(Min)	6.8 %	Vitamin A	(Min)	41,200 IU/lb
Calcium (Ca)	(Max)	8.2 %	Vitamin D-3	(Min)	4,100 IU/lb
Phosphorus (P)	(Min)	0.85 %	Vitamin E	(Min)	42 IU/lb
Salt (NaCl)	(Min)	2.7 %	Thiamin	(Min)	64 mg/lb

List of Ingredients

Soybean Meal, Calcium Carbonate, Distillers Dried Grains/Solubles, Pork Meat & Bone Meal, Cane Molasses, Salt, Linseed Meal, Wheat Middlings, Dehydrated Alfalfa Meal, Limestone ground, Lignin Sulfonate, Dried Aspergillus oryzae Fermentation Product, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Mineral Oil, Potassium Chloride, Sodium Selenite, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ethoxyquin Preservative, Thiamine Mononitrate.

Contains AMAFERM®, a registered trademark of BioZyme® Inc.

Feeding Directions

Mix 8 to 16 oz. per head, per day in Type C complete feed. Feed continuously to provide 0.25 to 0.50 mg of Melengestrol Acetate and 137.5-275 mg of lasalocid per head per day.

Caution

IMPORTANT: Total suppression of estrus is beyond the capabilities of melengestrol acetate due to individual heifer response to hormones, seasonal estrous cyclicity, management practices, and environmental stressors that may affect dry matter intake, thus melengestrol acetate daily dosage. Additive Cautions and Warnings: Not effective in steers and spayed heifers. The safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to supplements containing lasalocid as ingestion may be fatal. Mix thoroughly with grain and(or) roughage prior to feeding. Feeding undiluted, mixing errors and inadequate mixing of the total ration may result in excess lasalocid concentration which could be fatal to cattle. A withdrawal period has not been established for lasalocid in pre-ruminating cattle. Do not use in calves to be processed for veal. The feeding of supplemental selenium should not exceed 0.3 ppm of total diet. The addition to feed of higher levels of this premix containing selenium is not permitted.

Manufactured by:

BIOZYME®
INCORPORATED

MEDICATED

6010 Stockyards Expressway, St. Joseph, MO 64504
(800) 821-3070

Net Weight 50 lbs. (22.7 kg)

31 08/10/2016 145927E

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed. Safety Data Sheet available upon request.