

COXY CURB MED PEL Medicated

As an aid in the prevention or control of coccidiosis caused by Eimeria bovis and Eimeria zuernii.

ACTIVE DRUG INGREDIENT

| Amprolium | 1.25 % | | |
|-----------|---------------------|--|--|
| | Guarantood Analysis | | |

| | | Guaranteeu Analysis | | | | |
|----------------------|-------|---------------------|-------------|-------|--------|--|
| Crude Protein | (Min) | 11.2 % | Crude Fiber | (Max) | 17.9 % | |
| Crude Fat | (Min) | 3.1 % | ADF | (Max) | 19 % | |
| List of Tunue diauto | | | | | | |

List of Ingredients

Wheat Middlings, Soybean Hulls, Calcium Carbonate, Dehydrated Alfalfa Meal, Cane Molasses, Choice White Grease.

Feeding Directions

21 DAY PREVENTION PROGRAM: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, mix 0.04 pound of COXY CURB MED PEL Medicated into the feed for each 100 pounds of calf weight daily to provide 227 mg Amprolium per 100 lbs. bodyweight per day for 21 days.

| CALF WEIGHTS | COXY CUR lb/100 Head I | |
|-----------------|---------------------------|-----------|
| | Prevention | Treatment |
| 200 lbs. | 8 | 16 |
| 300 lbs. | 12 | 24 |
| 400 lbs. | 16 | 32 |
| 500 lbs. | 20 | 40 |
| 600 lbs. | 24 | 48 |
| | | |

5 DAY TREATMENT PROGRAM: When one or more calves show signs of coccidiosis, it is likely that the entire herd has been exposed. Treat on a herd basis by mixing 0.08 pound of COXY CURB MED PEL into the feed for each 100 pounds of calf weight daily to provide 454 mg Amprolium per 100 lbs. bodyweight per day for 5 days.

Caution

WARNING: Withdraw 24 hours before slaughter. Use as sole source of amprolium. A withdrawal period has not been established for this product in preruminating caves. Do not use in calves to be processed for veal.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS:

For prevention: Top-dress on or mix in the daily ration. Feed for 21 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard as the sole source of Amprolium. **For control**:Top-dress on or mix in the daily ration. Feed for 5 days as the sole source of Amprolium.

Manufactured by:



MEDICATED

6010 Stockyards Expressway, St. Joseph, MO 64504 (800) 821-3070 Net Weight 50 lbs. (22.7 kg) 31 08/21/2017 170500

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed. Safety Data Sheet available upon request.