



DECOXY MED PEL

Medicated

For the prevention of coccidiosis in ruminating and non-ruminating calves, including veal calves and cattle, caused by *Eimeria bovis* and *Eimeria zuernii*.

ACTIVE DRUG INGREDIENT

Decoquinatate 0.125 %

Guaranteed Analysis

| | | | | | |
|---------------|-------|--------|---------------|-------|-------|
| Crude Protein | (Min) | 14.1 % | Calcium (Ca) | (Min) | 7.6 % |
| Crude Fat | (Min) | 4.8 % | Calcium (Ca) | (Max) | 9.2 % |
| Crude Fiber | (Max) | 15.1 % | Potassium (K) | (Min) | 0.8 % |

List of Ingredients

Wheat Middlings, Distillers Dried Grains/Solubles, Soybean Hulls, Calcium Carbonate, Cane Molasses, Choice White Grease.

Feeding Directions

Feed at a rate to provide 22.7 mg Decoquinatate (.04 lb. or .64 oz.) per 100 lbs. of bodyweight daily. Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard.

FEEDING RATE OF DECOXY .125%

| BWT. | LB./H/D | OZ./H/D |
|------------|----------|----------|
| 100 lbs. | .04 lbs. | 0.64 oz. |
| 250 lbs. | .10 lbs. | 1.6 oz. |
| 500 lbs. | .20 lbs. | 3.2 oz. |
| 1,000 lbs. | .40 lbs. | 6.4 oz. |

Caution

WARNING: Do not feed to cows producing milk for food. Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard. Feed type C feed or milk replacer to provide 22.7 mg per 100 lb. bodyweight (0.5 mg/kg) per day.

CAUTION: Do not use in feeds containing bentonite.

Manufactured by:

BIOZYME®
INCORPORATED

MEDICATED

6010 Stockyards Expressway, St. Joseph, MO 64504
(800) 821-3070

Net Weight 50 lbs. (22.7 kg)

31 08/31/2017 170600

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed. Safety Data Sheet available upon request.